

# International Journal of Statistics and Applied Mathematics

ISSN: 2456-1452  
 Maths 2016; 1(1): 58-59  
 © 2016 Stats & Maths  
 www.mathsjournal.com  
 Received: 21-03-2016  
 Accepted: 24-04-2016

**MA Gopalan**  
 Professor, Department of  
 Mathematics, SIGC, Trichy,  
 Tamil Nadu. India

**S Vidhyalakshmi**  
 Professor, Department of  
 Mathematics, SIGC, Trichy,  
 Tamil Nadu. India

**R Presenna**  
 M. Phil Scholar, Department of  
 Mathematics, SIGC, Trichy,  
 Tamil Nadu. India

## Construction of special Dio-quadruple with property D (17)

**MA Gopalan, S Vidhyalakshmi and R Presenna**

### Abstract

This paper concerns with the study of constructing a special Dio quadruple (a, b, c, d) such that the product of any two elements of the set difference with their sum and increased by 17 is a perfect square.

**Keywords:** Diophantine Quadruples, Pell equation 2010 Mathematics subject classification 11D99

### Introduction

The problem of constructing the sets with property that the product of any two of its distinct elements is one less than a square has a very long history and such sets were studied by Diophantus [1]. A set of m positive integers  $\{a_1, a_2, \dots, a_m\}$  is said to have the property D (n),  $n \in \mathbb{Z} - \{0\}$  if  $a_i a_j + n$ , a perfect square for all  $1 \leq i < j \leq m$  and such a set is called a Diophantine m-tuples with property D(n). Many mathematicians considered the construction of different formulations of Diophantine quadruples with the property D (n) for any arbitrary integer n and also for any linear polynomials in n. In this context, one may refer [2-15] for an extensive review of various problems on Diophantine quadruples. This paper aims at constructing special dio – quadruple where the special mention is provided because it differs from the earlier one and the special Dio – quadruple is constructed where the product of any two members of the quadruple with the difference of the sum of the same members and the addition of seventeen satisfies the required property.

### Method of Analysis

Let  $a(k) = 4k + 5, b(k) = 4k - 3, k \geq 1$  be two integers such that  $a(k)b(k) - [a(k) + b(k)] + 17$  is a perfect square.

Let  $c_N(k)$  be any non-zero integer such that

$$a(k)c_N(k) - [a(k) + c_N(k)] + 17 = p_N^2(k)$$

$$b(k)c_N(k) - [b(k) + c_N(k)] + 17 = q_N^2(k)$$

Which, on simplification, results in the equations

$$(k + 1)c_N(k) - k + 3 = P_N^2(k); p_N(k) = 2P_N(k) \tag{1}$$

$$(k - 1)c_N(k) - k + 5 = Q_N^2(k); q_N(k) = 2Q_N(k) \tag{2}$$

Eliminating  $c_N(k)$  between (1) and (2), we obtain

$$(k - 1)p_N^2(k) - (k + 1)Q_N^2(k) = -8 \tag{3}$$

Taking  $P_N(k) = 2X_N(k) + (2k + 2)T_N(k) \tag{4}$

**Correspondence:**  
**MA Gopalan**  
 Professor, Department of  
 Mathematics, SIGC, Trichy,  
 Tamil Nadu. India

In (3), we have

$$X_N^2(k) = (k^2 - 1)T_N^2(k) + 1 \tag{6}$$

Which is satisfied by

$$X_N(k) = \frac{1}{2} \left[ (k + \sqrt{k^2 - 1})^{N+1} + (k - \sqrt{k^2 - 1})^{N+1} \right]$$

$$T_N(k) = \frac{1}{2\sqrt{k^2 - 1}} \left[ (k + \sqrt{k^2 - 1})^{N+1} - (k - \sqrt{k^2 - 1})^{N+1} \right] \tag{7}$$

Substituting N=0, 1, 2, 3,..... in (7) and using (4) & (1) in turn, we have

$$c_0(k) = 16k - 1$$

$$c_1(k) = 64k^3 - 16k + 1$$

$$c_2(k) = 256k^5 - 192k^3 + 32k + 1$$

$$c_3(k) = 1024k^7 - 1280k^5 + 448k^3 - 32k + 1 \text{ and so on.}$$

It is seen that the quadruple  $(a(k), b(k), c_{N-1}(k), c_N(k))$ , N=1, 2, 3,..... is a special dio-quadruple with property D(17).

A few illustrations are given below.

k	$(a(k), b(k), c_0(k), c_1(k))$	$(a(k), b(k), c_1(k), c_2(k))$	$(a(k), b(k), c_2(k), c_3(k))$
1	(9,1,17,49)	(9,1,49,97)	(9,1,97,161)
2	(13,5,33,481)	(13,5,481,6721)	(13,5,6721,93633)
3	(17,9,49,1681)	(17,9,1681,57121)	(17,9,57121,1940449)

**Conclusion**

This paper concerns with the construction of special dio - quadruples with property D (17). One may search for special dio-quadruples consisting of special numbers with suitable property.

**References**

1. Abu Muriefah FS, Al-Rashed A. some Diophantine quadruples in the ring  $Z[\sqrt{-2}]$  Math. Commun, 2004; 9:1-8.
2. Assaf E, Gueron S. Characterization of regular Diophantine quadruples, Elem. Math. 2001; 56:71-81.
3. Bashmakova IG. (ed.), Diophantus of Alexandria, Arithmetic's and the book of polynomial numbers, Nauka Moscow, 1974.
4. Dujella A. Some polynomial formulas for Diophantine quadruples, Grazer Math. Ber, 1996; 328:25-30.
5. Dujella A. Some estimates of the number of Diophantine quadruples, Publ. Math. Debrecen. 1998; 53:177-189.
6. Filipin A, Fujita Y. Any polynomial D (4)-quadruple is regular, Math. Commun, 2008; 13:45-55.
7. Filipin A. An irregular D (4)-quadruple cannot be extended to a quintuple, Acta Arith, 2009; 136:167-176.
8. Franusic Z. Diophantine quadruples in the ring  $Z[\sqrt{-2}]$  Math. Commun, 2004; 9:141-148.
9. Franusic Z. Diophantine quadruples in Quadratic Fields, Dissertation, University of Zagreb, 2005. (in Croatian)
10. Gopalan MA, Srividhya G. Diophantine quadruples for Fibonacci and Lucas Numbers with property D (4). diophantus J Math. 2012; 1:15-18.
11. Gopalan MA, Vidhyalakshmi S, Thiruniraiselvi N. A special Diophantine quadruple with property D (17), IJMTT, 2015; 20(2):110-112.
12. Gibbs P. Diophantine quadruples and Cayley's hyperdeterminant, XXX Mathematics Archive math. NT/0107203
13. Gibbs P. Adjugates of Diophantine Quadruples, Integers 2010; 10:201-209.
14. Srividhya G. Diophantine quadruples for Fibonacci numbers with property D (1), Indian journal of mathematics and Mathematical Sciences. 2009; 5:57-59.
15. Vidhyalakshmi S, Gopalan MA, Kavitha A. Special dio-quadruple with property D (5), IRJMEIT, 2014; 1(4):16-20.