

International Journal of Statistics and Applied Mathematics

ISSN: 2456-1452
Maths 2017; 2(2): 07-11
© 2017 Stats & Maths
www.mathsjournal.com
Received: 02-01-2017
Accepted: 03-02-2017

B SriRam

Department of Science and
Humanities, Acharya Nagarjuna
University College of
Engineering and Technology,
Acharya Nagarjuna University,
Guntur, India.

V Srinivas

Department of Science and
Humanities, Vijaya Institute of
Pharmaceutical Sciences for
Women, Vijayawada, India.

RRL Kantam

Departments of Statistics,
Acharya Nagarjuna University,
Guntur, India.

Correspondence

B SriRam

Department of Science and
Humanities, Acharya Nagarjuna
University College of
Engineering and Technology,
Acharya Nagarjuna University,
Guntur, India.

Significance test based on Rayleigh distribution

B SriRam, V Srinivas and RRL Kantam

Abstract

This paper deals with the Rayleigh distribution as a life time model. Moments of order statistics and an ordered sample are used to define a test statistic for the null hypothesis that the considered random variable has Rayleigh distribution. The percentiles of the test statistic are evaluated and the power of the test is computed. Comparison of Rayleigh distribution verses half-logistic distribution and gamma distribution with shape parameter 2 is presented.

Keywords: Testing, Moments, Order statistics, Power of Test

1. Introduction

The well-known Weibull distribution is one of the most widely used probability distribution in Reliability engineering discipline and also is studied by many researchers with respect to various problems on statistical inference. Weibull distribution with shape parameter 2 is known as Rayleigh distribution. In the field of statistical research, life testing experiments and reliability studies Rayleigh is the one of the most commonly used increasing failure rate models, whose probability density function is given by

$$f(x) = \frac{x}{\sigma^2} e^{-x^2/2\sigma^2}; x \geq 0 \quad (1.1)$$

$$= 0, \text{ otherwise}$$

where $\sigma > 0$, is the scale parameter of the distribution. The cumulative distribution function is given by

$$F(x) = 1 - e^{-x^2/2\sigma^2}; \text{ for } x \in [0, \infty) \quad (1.2)$$


$$= 0, \text{ otherwise}$$

2. The Graphs of frequency curve of Rayleigh distribution

The graph of frequency curve of Rayleigh distribution look similar to half logistic distribution, gamma distribution with shape parameter 2 models. Incidentally Rayleigh distribution is a combination of half logistic distribution and gamma distribution with shape parameter 2. We present below the frequency curve of Rayleigh distribution that look similar to half logistic distribution, gamma distribution with shape parameter 2.

In this paper we are interested to study the discrimination between Rayleigh distribution and half logistic distribution, gamma distribution with shape parameter 2. Another important of this study is to know whether half logistic distribution or gamma distribution with shape parameter 2 is a reasonable or better alternative to Rayleigh distribution in order to adopt the available simpler and admissible inferential procedures of half logistic distribution, gamma distribution with shape parameter 2 to Rayleigh distribution data. The similar studies of discrimination problems between probability models are made by Gupta *et al.* (2002) [6], Gupta and Kundu (2003A) [7], Gupta and Kundu (2003B) [8], Gupta and Kundu (2004) [9] Kundu and Manglick (2004) [12], Kundu *et al.* (2005) [14], Kundu and Manglick (2006) [11], Kundu (2005) and (2007) [14, 15], Kundu and Ragab (2005) [14], Arabin and Kundu (2009, 2010) [1, 2], Srinivasa Rao and Kantam (2011) [20], Kantam *et al.* (2015) [24] studied the significance test for the log-logistic distribution. Arabin and Kundu (2012A) [3], Arabin and Kundu (2012) [3], and the references there in.

Sultan (2007) ^[22] developed a test criterion to distinguish generalized exponential distribution from weibull, normal distributions, and moments of order statistics in samples drawn from generalized exponential distribution. In this paper we adopt the criterion suggested by Sultan (2007) ^[22] to distinguish between Rayleigh and half- logistic, gamma distribution with shape parameter 2. A brief description of procedure developed by Sultan (2007) ^[22] and its applications to our model is presented in Section 3. The methodology of arriving at the critical values and powers of the test procedures are given in Section 4 respectively. In all these cases, the percentiles of respective test statistic and powers of test procedures for selected sample sizes evaluated numerically and are tabulated in the respective situations. Rayleigh distribution is considered as null population (P₀) half-logistic distribution, gamma distribution with shape parameter 2 is considered as alternative populations (P₁).



3. Test of Significance

Let us assume that the distribution of the life of the product is the scaled Rayleigh distribution where probability density function and cumulative distribution function given by.

$$f_{x/\sigma}(x) = \frac{x}{\sigma^2} e^{-\frac{x^2}{2\sigma^2}} ; x>0, \sigma >0$$

$$= 0 \text{ otherwise}$$
(3.1)

$$F_{x/\sigma}(x) = 1 - e^{-\frac{x^2}{2\sigma^2}}$$

$$= 0 \text{ otherwise}$$
(3.2)

And its probability density function of standard Rayleigh distribution as given by

$$f(z) = \frac{z}{\sigma} e^{-\frac{z^2}{2\sigma^2}} ; z \geq 0$$

$$= 0 \text{ otherwise}$$
(3.3)

$$F(z) = 1 - e^{-\frac{z^2}{2\sigma^2}} \text{ for } z \in [0, \infty)$$

$$= 0 \text{ otherwise}$$
(3.4)

Probability density function of scaled half logistic distribution as given by

$$f_{x/\sigma}(x) = \frac{2e^{-\frac{x}{\sigma}}}{(1 + e^{-\frac{x}{\sigma}})^2} ; x \geq 0, \sigma > 0$$

$$= 0 \text{ otherwise}$$
(3.5)

$$F_{x/\sigma}(x) = \frac{1 - e^{-\frac{x}{\sigma}}}{1 + e^{-\frac{x}{\sigma}}}; x \geq 0, \sigma > 0$$

$$= 0 \text{ otherwise} \tag{3.6}$$

Probability density function of standard half- logistic distribution as given by

$$f(z) = \frac{2e^{-z}}{(1 + e^{-z})^2}; z \geq 0$$

$$= 0 \text{ otherwise} \tag{3.7}$$

and its cumulative distributive function

$$F(z) = \frac{1 - e^{-z}}{1 + e^{-z}}; z \geq 0$$

$$= 0 \text{ otherwise} \tag{3.8}$$

Probability density function of scaled gamma distribution with shape parameter 2 is given by

$$f_{x/\sigma}(x) = \frac{x}{\sigma^2} e^{-\frac{x}{\sigma}}; x > 0, \sigma > 0$$

$$= 0 \text{ otherwise} \tag{3.9}$$

and its cumulative distributive function

$$F_{x/\sigma}(x) = (1 - e^{-\frac{x}{\sigma}})(1 + \frac{x}{\sigma})$$

$$= 0 \text{ otherwise} \tag{3.10}$$

Probability density function of standard gamma distribution with shape parameter 2 is given by

$$f(z) = z \cdot e^{-z}; z \geq 0$$

$$= 0 \text{ otherwise} \tag{3.11}$$

and its cumulative distributive function of standard gamma distribution with shape parameter 2 is given by

$$F(z) = (1 - e^{-z})(1 + z)$$

$$= 0 \text{ otherwise} \tag{3.12}$$

Let x_1, x_2, \dots, x_n be a random sample of size n , here we test the null hypothesis P_0 : the sample has come from Rayleigh distribution against each of the alternative hypothesis against the one of the alternative hypothesis.

1. P_1 : the sample has come from half logistic distribution.
2. P_1 : the sample has come from gamma distribution with shape parameter 2.

As we mentioned in Section 1 that Rayleigh distribution is a combination of half- logistic distribution or gamma distribution with shape parameter 2. From this context we have considered Rayleigh distribution as null population and half-logistic distribution, gamma distribution with shape parameter 2 as alternative population. Sultan (2007) suggested a test statistic given by a formula

$$T = \frac{\sum_{i=0}^n X_{(i)} \alpha_{(i)}}{\sqrt{\sum_{i=0}^n X_{(i)}^2 \sum_{i=0}^n \alpha_{(i)}^2}} \tag{3.13}$$

where $x_{(i)}$ is the order observation in the sample, $\alpha_{(i)}$ is the expected value of i^{th} standard order statistic in a sample of size n from the null population. If x_1, x_2, \dots, x_n is a sample of size n from the null population the Equation (3.13) for T is used as a test statistic to discriminate a null population and the corresponding alternative population with the help of the critical values. Hence, the sampling distribution of T and its percentiles therefore are essential to make use of the test statistic T . Since, sample quantiles are consistent estimators of population quantiles. Hence we propose a statistic similar to T based on population quantiles rather than moments of order statistics and is given by

$$T^* = \frac{\sum_{i=1}^n \delta_{(i)} x_{(i)}}{\sqrt{\sum_{i=1}^n x_{(i)}^2 \sum_{i=1}^n \delta_{(i)}^2}} \tag{3.14}$$

where $\delta_{(i)} = F^{-1}(p) = \left(\frac{p_i}{1-p_i}\right)^{\frac{1}{\beta}}$ with $p_i = \frac{i}{n+1}$

We have tabulated the percentiles of empirical sampling distribution of T^* for $n=5(5)25$ through 10,000 Monte-Carlo simulation runs and are given in Table 3.1. The percentiles of T^* would serve as critical values to test null hypothesis that a given sample comes from Rayleigh distribution and half logistic distribution, gamma distribution with shape parameter 2. Therefore a large

value of T^* implies a strong linear relation between $x_{(i)}$ and $\delta_{(i)}$, which means that the sample is taken from Rayleigh distribution. On the other hand, a small value of T^* indicates that the sample is from a distribution different from Rayleigh distribution. Hence the statistic T^* is given by Equation (3.14) is approximate for testing the null hypothesis that X is distributed according to a Rayleigh distribution.

The decision to reject the null hypothesis is based on the critical values of T with a pre-assigned level of significance. Accordingly, the percentiles of T^* are essential to carry out the test procedure regarding the above hypothesis. For example, let t_α denote the α -percentile of T^* , then if

$$P_T = P\left\{\left(\frac{t}{t} > t_\alpha\right)\right\} = 1 - \alpha \tag{3.15}$$

Holds, the null hypothesis cannot be rejected with level α . Thus, the percentiles of T^* are essential, but unfortunately it is not possible to obtain the distribution of T^* analytically. We determined the percentiles of T^* by means of simulation on the basis of 10,000 simulation runs for various sample sizes. The obtained percentiles are given in Table 3.2.

4. Power of the test

The power of a test refers to the probability of not rejecting a wrong null hypothesis and this probability depends heavily on the assumed alternative distribution. Here in this case, the null hypothesis given by Rayleigh distribution as the alternative distribution we consider half- logistic distribution. The test statistic T^* is calculated by generating a sample of a specified size from each of the alternative population with the $\delta_{(i)}$ in the formula for T^* being those of Rayleigh distribution. The corresponding calculated value of T^* is compared with the corresponding critical value given in Table 4.1. This process is repeated 10,000 times and the number of rejections of null hypothesis out of 10,000 times is taken as a measure of power of the test i.e.,

$$\text{power of the test} = \frac{\text{number of rejections of } H_0}{10,000} \tag{4.1}$$

The calculation of the power were performed for the above mentioned alternative with various sample sizes and various levels of significance for complete samples were presented in Tables 4.1 and 4.2 and the tables indicate the high power of the proposed test for all considered alternatives. The power indicates that the half logistic alternative is more powerful and gamma distribution with shape parameter 2 is less sensitive as alternative population. This shows that the test statistic clearly discriminates a Rayleigh distribution from half logistic distribution and gamma distribution with shape parameter 2 with decreasing power in that order.

Table 3.1: Percentiles of T^* for complete samples

$\frac{N}{P}$	5	10	15	20	25
0.00135	0.867997	0.914191	0.99118	0.999216	0.99927
0.005	0.897317	0.938334	0.998805	0.998957	0.999082
0.01	0.919219	0.94823	0.98864	0.998782	0.998915
0.025	0.93836	0.961392	0.998315	0.998497	0.998691
0.05	0.952971	0.969693	0.99739	0.997915	0.998428
0.10	0.965421	0.977381	0.99739	0.997767	0.998058
0.90	0.99637	0.996767	0.983222	0.986755	0.988884
0.95	0.997511	0.997486	0.977828	0.982839	0.985089
0.975	0.998266	0.998019	0.971573	0.978253	0.9881579
0.99	0.998896	0.998433	0.962629	0.975645	0.975645
0.995	0.999311	0.998749	0.955778	0.9764860	0.970486
0.99865	0.999508	0.999105	0.941952	0.962572	0.962572

Table 3.2: Estimates of the power of test for two alternative distributions

N	Half-logistic distribution			Gamma(2)		
	0.99	0.95	0.90	0.99	0.95	0.90
5	0.9665	0.9823	0.9665	0.9955	0.9748	0.9428
10	0.9986	0.9947	0.9903	0.9962	0.983	0.9662
15	0.9997	0.998	0.9957	0.9975	0.9879	0.9746
20	0.9997	0.9991	0.9977	0.9982	0.9905	0.9825
25	1	0.9999	0.9995	0.9989	0.994	0.9861

Table 4.1: Power T values for Rayleigh vs HLD

S.No	Sample no	99%	95%	90%
1	5	0.0335	0.0177	0.0020
2	10	0.0097	0.0053	0.0014
3	15	0.0043	0.0020	0.0003
4	20	0.0003	0.0009	0.0003
5	25	0.0005	0.0001	0

Table 4.2: Power T values for Rayleigh vs Gamma (2)

S.No	Sample no	99%	95%	90%
1	5	0.0572	0.0252	0.0045
2	10	0.0338	0.0170	0.0038
3	15	0.0254	0.0121	0.0025
4	20	0.0175	0.0095	0.0018
5	25	0.0139	0.0060	0.0013

References

1. Arabin K.D, Kundu D. Discriminating among the log-normal, weibull and generalized exponential distributions, *IEEE Transactions on Reliability* 2009; 58:416-424.
2. Arabin K.D, Kundu D. Discriminating between the log-normal and log-logistic distributions, *Communications in Statistics - Theory and Methods* 2010; 39:280-292.
3. Arabin K.D, Kundu D. Discriminating between the weibull and log-normal distributions for type-II censored data, *Statistics* 2012a; 46:197-214.
4. Arabin K.D, Kundu D. Discriminating between bivariate generalized exponential and bivariate weibull distributions, *Chilean Journal of Statistics*. 2012b; 3:93-110.
5. Block H.W, Rao. A Beta warning-time distribution and a distended beta distribution, *Sankhya B* 1973; 35:79-84.
6. Gupta R.D, Kundu D. Generalized exponential distributions: statistical inferences, *Journal of Statistical Theory and Applications*. 2002; 1:101-118.
7. Gupta R.D, Kundu D. Discriminating between the weibull and the generalized exponential distributions, *Computational Statistics and Data Analysis* 2003a; 43:179-196.
8. Gupta R.D, Kundu D. Closeness of gamma and generalized exponential distribution, *Communications in Statistics - Theory and Methods* 2003b; 32:705-721.
9. Gupta R.D, Kundu D. Discriminating between gamma and generalized exponential distributions, *Journal of Statistical Computation and Simulation*. 2004; 74:107-121.
10. Kundu D, Gupta R.D, Manglick A. Discriminating between the log-normal and the generalized exponential distributions, *Journal of the Statistical Planning and Inference*. 2005; 127:213-227.
11. Kundu D, Gupta R.D, Manglick A. A convenient way of generating normal random variables using generalized exponential distribution, *Journal of Modern Applied Statistical Methods*. 2006; 5:300-306.
12. Kundu D, Manglick A. Discriminating between the weibull and log-normal distributions, *Naval Research Logistics* 2004; 51:893-905.
13. Kundu D, Manglick A. Discriminating between the gamma and log-normal distributions, *Journal of Applied Statistical Sciences* 2005; 14:175-187.
14. Kundu D, Raqab M.Z. Generalized Rayleigh Distribution: Different methods of estimation, *Computational Statistics and Data Analysis* 2005; 49:187-200.
15. Kundu D, Raqab M.Z. Discriminating between the log-normal and generalized rayleigh distributions, *Statistics* 2007; 41:505-515.
16. Malik H.J. Exact distribution of the quotient of independent generalized gamma variables, *Canadian Mathematical Bulletin* 1967; 10:463-466.
17. Ragab A, Green J. On order statistics from the log-logistic distribution and their properties, *Communications in Statistics - Theory and Methods* 1984; 13:2713-2724.
18. Ragab A, Green J. Estimation of the parameters of the log-logistic distribution based on order statistics, *American Journal of Mathematical and Management Science*. 1987; 7:307-324
19. Shah B.K, Dave P.H. A note on log-logistic distribution, *Journal of the M.S. University of Baroda (Science Number)* 1963; 12:21-22.
20. Srinivasa Rao B, Kantam R.R.L. Significance test for half logistic distribution, *Economic Quality Control* 2011; 26:53-62.
21. Srinivasa Rao G, Kantam R.R.L. Two stage estimation in log-logistic model, *Pakistan Journal of Statistics*. 2012c; 28:331-339.
22. Sultan K.S. Order Statistics from the Generalised exponential distribution and applications, *Communications in Statistics - Theory and Methods* 2007; 36:1409-1418.
23. Tadikamalla P.R, Johnson N.L. Systems of frequency curves generated by the transformation of logistic variables, *Biometrika* 1982; 69:461-465.
24. Kantam R.R.L, SriRam B, Suhasini. Significance test based on Log-logistic distribution, *International Journal of computational and Theoretical statistics*, ISSN (2210-1519) Int. J Comp Tho Stat No. 2, Nov, 2015)