

International Journal of Statistics and Applied Mathematics



ISSN: 2456-1452
 Maths 2017; 2(6): 155-158
 © 2017 Stats & Maths
 www.mathsjournal.com
 Received: 04-09-2017
 Accepted: 05-10-2017

Pramod Shinde
 Research Scholar
 Opjs University, Churu,
 Rajasthan, India

Dr. Ashwini Nagpal
 Associate Professor
 Department of Mathematics
 Opjs University, Churu,
 Rajasthan, India

Multiple integration of a function

Pramod Shinde and Dr. Ashwini Nagpal

Abstract

The multiple integral is a generalization of the definite integral to functions of more than one real variable, for example, $f(x, y)$ or $f(x, y, z)$. Integrals of a function of two variables over a region in R^2 are called double integrals, and integrals of a function of three variables over a region of R^3 are called triple integrals.

Just as the definite integral of a positive function of one variable represents the area of the region between the graph of the function and the x-axis, the double integral of a positive function of two variables represents the volume of the region between the surface defined by the function (on the three-dimensional Cartesian plane where $z = f(x, y)$) and the plane which contains its domain. (The same volume can be obtained via the triple integral—the integral of a function in three variables—of the constant function $f(x, y, z) = 1$ over the above-mentioned region between the surface and the plane.) If there are more variables, a multiple integral will yield hypervolumes of multidimensional functions.

Keywords: multiple integration, function, variable

Introduction

Multiple integration of a function in n variables: $f(x_1, x_2, \dots, x_n)$ over a domain D is most commonly represented by nested integral signs in the reverse order of execution (the leftmost integral sign is computed last), followed by the function and integrand arguments in proper order (the integral with respect to the rightmost argument is computed last).

Multiple integrals have many properties common to those of integrals of functions of one variable (linearity, commutativity, monotonicity, and so on). One important property of multiple integrals is that the value of an integral is independent of the order of integrands under certain conditions. This property is popularly known as Fubini's theorem

When the domain of integration is symmetric about the origin with respect to at least one of the variables of integration and the integrand is odd with respect to this variable, the integral is equal to zero, as the integrals over the two halves of the domain have the same absolute value but opposite signs. When the integrand is even with respect to this variable, the integral is equal to twice the integral over one half of the domain, as the integrals over the two halves of the domain are equal.

The integral of order ν of a function $f(r)$ is given by:

$$F_\nu(k) = \int_0^\infty f(r) J_\nu(kr) r dr$$

where

is J_ν the Bessel function of the first kind of order ν with $\nu \geq -\frac{1}{2}$

The inverse integral of $F_\nu(k)$ is defined as:

$$f(r) = \int_0^\infty F_\nu(k) J_\nu(kr) k dk$$

which can be readily verified using the orthogonality relationship described below.

Correspondence
Pramod Shinde
 Research Scholar
 Opjs University, Churu,
 Rajasthan, India

Inverting a integral of a function $f(r)$ is valid at every point at which $f(r)$ is continuous provided that the function is defined in $(0, \infty)$, is piecewise continuous and of bounded variation in every finite subinterval in $(0, \infty)$, and

$$\int_0^\infty |f(r)| r^{\frac{1}{2}} dr < \infty.$$

However, like the integral, the domain can be extended by a density argument to include some functions whose above integral is not finite, for example

$$f(r) = (1 + r)^{-3/2}$$

An alternative definition says that the integral of $g(r)$ is:

$$h_\nu(k) = \int_0^\infty g(r) J_\nu(kr) \sqrt{kr} dr$$

The two definitions are related:

$$\text{If } g(r) = f(r)\sqrt{r} \text{ then } h_\nu(k) = F_\nu(k)\sqrt{k}.$$

This means that, as with the previous definition, the integral defined this way is also its own inverse:

$$g(r) = \int_0^\infty h_\nu(k) J_\nu(kr) \sqrt{kr} dk$$

The obvious domain now has the condition

$$\int_0^\infty |g(r)| dr < \infty$$

but this can be extended.

According to the reference given above, we can take the integral as the limit as the upper limit goes to infinity (an improper integral rather than a Lebesgue integral) and in this way the integral and its inverse work for all functions in $L^2(0, \infty)$.

The Bessel functions form an orthogonal basis with respect to the weighting factor r :

$$\int_0^\infty J_\nu(kr) J_\nu(k'r) r dr = \frac{\delta(k - k')}{k}, \quad k, k' > 0.$$

Research Study

If $f(r)$ and $g(r)$ are such that their integral $F_\nu(k)$ and $G_\nu(k)$ are well defined, then the Plancherel theorem states

$$\int_0^\infty f(r)g(r)r dr = \int_0^\infty F_\nu(k)G_\nu(k)k dk.$$

Parseval's theorem, which states:

$$\int_0^\infty |f(r)|^2 r dr = \int_0^\infty |F_\nu(k)|^2 k dk,$$

is a special case of the Plancherel theorem. These theorems can be proven using the orthogonality property.

The integral of order zero is essentially the 2-dimensional Fourier transform of a circularly symmetric function.

Consider a 2-dimensional function $f(\mathbf{r})$ of the radius vector \mathbf{r} . Its Fourier transform is:

$$F(\mathbf{k}) = \iint f(\mathbf{r})e^{i\mathbf{k}\cdot\mathbf{r}} d\mathbf{r}.$$

With no loss of generality, we can pick a polar coordinate system (r, θ) such that the k vector lies on the $\theta = 0$ axis (in K -space). The Fourier transform is now written in these polar coordinates as:

$$F(\mathbf{k}) = \int_{r=0}^{\infty} \int_{\theta=0}^{2\pi} f(r, \theta) e^{i\mathbf{k}\cdot\mathbf{r}} r \, d\theta \, dr$$

where θ is the angle between the \mathbf{k} and \mathbf{r} vectors. If the function f happens to be circularly symmetric, it will have no dependence on the angular variable θ and may be written $f(r)$. The integration over θ may be carried out, and the Fourier transform is now written:

$$F(\mathbf{k}) = F(k) = 2\pi \int_0^{\infty} f(r) J_0(kr) r \, dr$$

which is just 2π times the zero-order integral of $f(r)$. For the reverse transform,

$$f(r) = \frac{1}{(2\pi)^2} \iint F(\mathbf{k}) e^{-i\mathbf{k}\cdot\mathbf{r}} \, d\mathbf{k} = \frac{1}{2\pi} \int_0^{\infty} F(k) J_0(kr) k \, dk$$

so $f(r)$ is $1/2\pi$ times the zero-order integral of $F(k)$.

Significance of the Study

For an n -dimensional Fourier transform,

$$F(\mathbf{k}) = \int f(\mathbf{r}) e^{-i\mathbf{k}\cdot\mathbf{r}} \, d^n \mathbf{r}$$

if the function f is radially symmetric, then:

$$k^{n/2-1} F(k) = (2\pi)^{n/2} \int_0^{\infty} r^{n/2-1} f(r) J_{n/2-1}(kr) r \, dr$$

To generalize: If f can be expanded in a multipole series,

$$f(r, \theta) = \sum_{m=-\infty}^{\infty} f_m(r) e^{im\theta},$$

and if θ_k is the angle between the direction of \mathbf{k} and the $\theta = 0$ axis,

$$\begin{aligned} F(\mathbf{k}) &= \int_0^{\infty} r \, dr \int_0^{2\pi} d\theta f(r, \theta) e^{i\mathbf{k}\cdot\mathbf{r}} \\ &= \sum_m \int_0^{\infty} r \, dr \int_0^{2\pi} d\theta f_m(r) e^{im\theta} e^{i\mathbf{k}\cdot\mathbf{r}} \\ &= \sum_m e^{im\theta_k} \int_0^{\infty} r \, dr f_m(r) \int_0^{2\pi} d\varphi e^{im\varphi} e^{i\mathbf{k}\cdot\mathbf{r}} \quad \varphi = \theta - \theta_k \\ &= \sum_m e^{im\theta_k} \int_0^{\infty} r \, dr f_m(r) 2\pi i^m J_m(kr) \\ &= 2\pi \sum_m i^m e^{im\theta_k} \int_0^{\infty} f_m(r) J_m(kr) r \, dr. \\ &= 2\pi \sum_m i^m e^{im\theta_k} F_m(k) \end{aligned}$$

where $F_m(k)$ is the m -th order Hankel transform of $f_m(r)$.

Additionally, if f_m is sufficiently smooth near the origin and zero outside a radius R , it may be expanded into a Chebyshev series,

$$f_m(r) = r^m \sum_{t \geq 0} f_{mt} \left(1 - \left(\frac{r}{R}\right)^2\right)^t, \quad 0 \leq r \leq R.$$

So that

$$\begin{aligned} F(\mathbf{k}) &= 2\pi \sum_m \tilde{r}^m e^{im\theta_k} \sum_t f_{mt} \int_0^R r^m \left(1 - \left(\frac{r}{R}\right)^2\right)^t J_m(kr) r \, dr \quad (*) \\ &= 2\pi \sum_m \tilde{r}^m e^{im\theta_k} R^{m+2} \sum_t f_{mt} \int_0^1 x^m (1-x^2)^t J_m(kxR) x \, dx \quad x = \frac{r}{R} \\ &= 2\pi \sum_m \tilde{r}^m e^{im\theta_k} R^{m+2} \sum_t f_{mt} \frac{t! 2^t}{(kR)^{1+t}} J_{m+t+1}(kR). \end{aligned}$$

The above can be viewed as a more general case that is not as constrained as the previous case in the previous section. The numerically important aspect is that the expansion coefficients f_{mt} are accessible with Discrete Fourier transform techniques. Insertion into the previous formula yields.

References

1. Gaskill Jack D. Linear Systems, Fourier Transforms, and Optics. New York: John Wiley & Sons, 2008. ISBN 0-471-29288-5.
2. Polyanin AD, Manzhirov AV. Handbook of Integral Equations. Boca Raton: CRC Press, 2008. ISBN 0-8493-2876-4.
3. Smythe, William R. Static and Dynamic Electricity (3rd ed.). New York: McGraw-Hill. 2008, 179-223.
4. Offord AC. On integral. Proceedings of the London Mathematical Society. 2012; 39(2):49-67.
5. Eason G, Noble B, Sneddon IN. On certain integrals of Lipschitz-Hankel type involving products of Bessel Functions. Philosophical Transactions of the Royal Society. 2005; 247(935):529-551.
6. Kilpatrick JE, Katsura Shigetoshi, Inoue Yuji. Calculation of integrals of products of Bessel functions. Mathematics of Computation. 2007; 21(99):407-412.
7. MacKinnon, Robert F. The asymptotic expansions of Hankel transforms and related integrals. Mathematics of Computation. 2012; 26(118):515-527.
8. Linz Peter, Kropp TE. A note on the computation of integrals involving products of trigonometric and Bessel functions. Mathematics of Computation. 2013; 27(124):871-872.
9. Noll, Robert J. Zernike polynomials and atmospheric turbulence. Journal of the Optical Society of America. 2009; 66(3):207-211.
10. Siegman AE. Quasi-fast Hankel transform. Opt. Lett. 2010; 1(1):13-15.
11. Magni Vittorio, Cerullo Giulio, De Silverstri Sandro. High-accuracy fast Hankel transform for optical beam propagation. J Opt. Soc. Am. 2012; 9(11):2031-2033.
12. Agnesi A, Reali Giancarlo C, Patrini G, Tomaselli A. Numerical evaluation of the Hankel transform: remarks. Journal of the Optical Society of America. 2013; 10(9).
13. Barakat Richard. Numerical evaluation of the zero-order Hankel transform using Filon quadrature philosophy. Applied Mathematics Letters. 2006; 9(5):21-26.