

International Journal of Statistics and Applied Mathematics

ISSN: 2456-1452
 Maths 2018; 3(5): 62-68
 © 2018 Stats & Maths
 www.mathsjournal.com
 Received: 19-07-2018
 Accepted: 20-08-2018

Akintunde Oyetunde A
 Department of Mathematics,
 Faculty of Science, Federal
 University Oye-Ekiti,
 Ekiti State, Nigeria

Statistical hybridization of normal and Weibull distributions with its properties and applications

Akintunde Oyetunde A

Abstract

The normal distribution is one of the most popular probability distributions with applications to real life data. In this research paper, an extension of this distribution together with Weibull distribution called the Weimal distribution which is believed to provide greater flexibility to model scenarios involving skewed data was proposed. The probability density function and cumulative distribution function of the new distribution can be represented as a linear combination of exponential normal density functions. Analytical expressions for some mathematical quantities comprising of moments, moment generating function, characteristic function and order statistics were presented. The estimation of the proposed distribution's parameters was undertaken using the method of maximum likelihood estimation. Two data sets were used for illustration and performance evaluation of the proposed model. The results of the comparative analysis to other baseline models show that the proposed distribution would be more appropriate when dealing with skewed data.

Keywords: Weibull-Generalized family of distributions, maximum likelihood, moment generating function, characteristic function, order statistics

1. Introduction

One of the main motivations for studying new families of statistical distributions lies in the increased flexibility of fitting various datasets that cannot be properly fitted by existing distributions. In many applied areas, such as environmental and medical sciences, engineering, biological studies, lifetime analysis, actuaries, economics, as well as finance and insurance; there is a clear need for extended forms of these distributions. The normal distribution is the most popular probability model having wider applications in solving real life problems. When the number of observations is large, it can serve as an approximation to other probability models.

The normal distribution is also called Guassian distribution, named after the German Mathematician Carl Freidrich Gauss (1777-1855) who introduced it in connection with the theory of error. The probability density function (pdf) and the cumulative distribution function (cdf) of the normal distribution with location parameter $-\infty < \mu < \infty$ and scale parameter $\sigma > 0$ are given respectively by:

$$f(x, \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) = \frac{1}{\sigma} \phi\left(\frac{x-\mu}{\sigma}\right) \quad (1)$$

$$F(x, \mu, \sigma) = \Phi\left(\frac{x-\mu}{\sigma}\right) \quad (2)$$

Attempts to generalize the Normal distribution have led to the development of Skewed Normal distribution [1], the Beta-Normal distribution [2], the Generalized Normal distribution [3], the Kumaraswamy-Normal distribution [4], the McDonald-Normal distribution [5], the modified Beta-Normal distribution [6], the Gamma-Normal distribution [7], the modified Gamma-Normal distribution [8], the Kummer Beta-Normal distribution [9], and a host of others. These distributions are proven to be more flexible than the classical Normal distribution when applied to the real life datasets.

Correspondence
Akintunde Oyetunde A
 Department of Mathematics,
 Faculty of Science, Federal
 University Oye-Ekiti,
 Ekiti State, Nigeria

Several generalized families of distributions have been proposed in the literature. For instance, the Beta-G [2], the Kuraraswamy-G [4], Transmuted family of distributions [10, Gamma-G (type 1) [11], McDonald-G [12], Gamma-G (type 2) [13], Gamma-G (type 3) [14], Log-Gamma-G [15], Exponentiated T-X [16], Logistic-G [18], Gamma-X and Weibull-X [19], Logistic-X [20], Weibull-G [21], Beta Marshall-Oikin family of distribution [22], and many others are available in literature. The focus of this research article is the Weibull-G family of distributions [21] which can be obtained by using the odds ratio of failure rate because they highlighted the Normal as one of the distribution that can be obtained by their Weibull generator.

The use of four-parameter distribution should be sufficient for most practical purposes, and at least three-parameter are needed in such distributions as stated in [23], but doubted any noticeable improvement arising from including a fifth- or sixth parameter. The Weibull-G family of distributions have been adopted by several notable researchers among to generate known theoretical models such as the Weibull-Exponential distribution [24], Weibull-rayleigh distribution [25], and Weibull-Frechet distribution [26]. In this research paper, a proposition of a probability model called a Weibull-Normal distribution, also to be known as Weimal distribution, which resulted from hybridizing Weibull and Normal distributions by utilizing the Weibull-G family generator proposed in [21].

2. The Weibull-Generalized (Weibull-G) family of Distribution

Given any continuous baseline distribution with *cdf* $G(x, \xi)$ and *pdf* $g(x, \xi)$ aiming at providing greater flexibility in modeling real-life datasets; a Weibull generalized family of distribution, according to [21], will have a *cdf* $G(x, \xi)$ defined by:

$$F_{WG}(x, \alpha, \beta, \xi) = \int_{-\infty}^{G(x, \xi)} \alpha \beta t^{\beta-1} e^{-at^\beta} dt \tag{3}$$

The above integral yields:

$$1 - \exp \left\{ -\alpha \left[\frac{G(x, \xi)}{\bar{G}(x, \xi)} \right]^\beta \right\}, x \in D \subseteq \mathbb{R}, \alpha, \beta > 0 \tag{4}$$

The corresponding *pdf* is given by:

$$F_{WG}(x, \alpha, \beta, \xi) = \alpha \beta g(x, \xi) \frac{G(x, \xi)^{\beta-1}}{\bar{G}(x, \xi)^{\beta+1}} \exp \left\{ -\alpha \left[\frac{G(x, \xi)}{\bar{G}(x, \xi)} \right]^\beta \right\} \tag{5}$$

Where $g(x, \xi)$ and $G(x, \xi)$ are the respective *pdf* and *cdf* of the baseline distribution indexed by parameter vector ξ , where $\alpha > 0$ and $\beta > 0$ are the scale and shape parameters respectively.

Equation (5) gives the *pdf* of any Weibull-G family of distribution and is most tractable when both *cdf* and *pdf* have simple analytic expressions. The major benefit of the Weibull-generator expressed in equation (3) lies in its ability to offer more flexibility to the extremes of the *pdf* and this makes it more suitable for analyzing data with high degree of asymmetry.

2.1 The Weibull-Normal distribution

Taking into account the *pdf* and *cdf* of the normal distribution as given in equations (1) and (2) with location parameter $\mu \in \mathbb{R}$ and dispersion parameter $\alpha > 0$. The respective *cdf* and *pdf* of the proposed four-parameter Weibull-Normal distribution can be obtained from equations (4) and (5) as follows:

$$F(x) = F(x, \alpha, \beta, \mu, \sigma) = 1 - \exp \left\{ -\alpha \left[\frac{\phi\left(\frac{x-\mu}{\sigma}\right)}{1-\phi\left(\frac{x-\mu}{\sigma}\right)} \right]^\beta \right\} \tag{6}$$

And

$$f(x) = f(x, \alpha, \beta, \mu, \sigma) = \frac{\alpha \beta}{\sigma} \phi\left(\frac{x-\mu}{\sigma}\right) \frac{\phi\left(\frac{x-\mu}{\sigma}\right)^{\beta-1}}{\left(1-\phi\left(\frac{x-\mu}{\sigma}\right)\right)^{\beta+1}} \exp \left\{ -\alpha \left[\frac{\phi\left(\frac{x-\mu}{\sigma}\right)}{1-\phi\left(\frac{x-\mu}{\sigma}\right)} \right]^\beta \right\} \tag{7}$$

The plots of the pdf and cdf of the new Weibull-Normal distribution for the selected parameter values and plotted and it was observed that the cdf of the Weibull-Normal distribution increases as x increases and approaches one as x gets larger. The different pdf plots of the Weimal distribution under different parameter values indicate that it is negatively skewed distribution and hence it will be very appropriate in modeling skewed real-life datasets unlike the symmetric normal distribution. It is interesting to note that whenever $\alpha=\beta=1$, then the Weimal distribution becomes normal and that all parameter values affect the graph of the pdf in different directions and different rates.

2.2 Useful Extensions

Extensions for equations (6) and (7) can be derived using the concept of exponentiated distributions as follows: Consider the exponentiated normal (EN) distribution as stated by Nadarajah and Kotz [6] with power parameter $a > 0$ defined by $Y \sim EN(a, \mu, \sigma)$ with *cdf* and *pdf* given respectively by:

$$H_a(y) = H(y, a) = \phi\left(\frac{y-\mu}{\sigma}\right)^a \tag{8}$$

And

$$h_a(y, a) = h(y, a) = \frac{a}{b} \phi\left(\frac{x-\mu}{\sigma}\right) \phi\left(\frac{x-\mu}{\sigma}\right)^{a-1} \tag{9}$$

By expanding the exponential term in equation (7) using power series and utilizing the generalized binomial theorem while substituting, we can re-write equation (7) as:

$$f(x) = \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \frac{(-1)^k a^{k+1} \Gamma(\beta(k+1)+j+1)}{j! k! \sigma \Gamma(\beta(k+1)+1)} \phi\left(\frac{x-\mu}{\sigma}\right) \phi\left(\frac{x-\mu}{\sigma}\right)^{\beta(k+1)+j-1} \tag{10}$$

Using the result in equation (8), we can now express the *pdf* of the Weimal distribution as a linear combination of exponentiated (exponential-G) density functions as:

$$f(x) = \sum_{j,k=0}^{\infty} w_{j,k} h_{\beta(k+1)+j}(x) \tag{11}$$

Where

$$h_{\beta(k+1)+j}(x) = h(x; h_{\beta(k+1)+j}, \mu, \sigma) = [\beta(k+1) + j] \phi\left(\frac{x-\mu}{\sigma}\right) \phi\left(\frac{x-\mu}{\sigma}\right)^{\beta(k+1)+j-1}$$

Corresponding to:

$$h_a(x) = h(x, a, \mu, \sigma) = a \phi\left(\frac{x-\mu}{\sigma}\right) \phi\left(\frac{x-\mu}{\sigma}\right)^{a-1}$$

Denotes the exponential normal *pdf* ($X \sim EN(\beta(k+1) + j, \mu, \sigma)$) and the coefficient of the distribution $w_{j,k}$ is given by:

$$w_{j,k} = \frac{(-1)^k a^{k+1} \beta \Gamma(\beta(k+1) + j + 1)}{j! k! \sigma [\beta(k+1) + j] \Gamma(\beta(k+1) + 1)}$$

By integrating the *pdf* in equation (9) with respect to x , we obtain the corresponding *cdf* as:

$$F(x) = \sum_{j,k=0}^{\infty} w_{j,k} H(x; h_{\beta(k+1)+j}, \mu, \sigma) = \sum_{j,k=0}^{\infty} w_{j,k} \phi\left(\frac{x-\mu}{\sigma}\right)^{\beta(k+1)+j} \tag{12}$$

If $\beta > 0$ is a real number (positive non-integer), we can expand the last term in (12) as:

$$\left(\frac{x-\mu}{\sigma}\right)^{\beta(k+1)+j} = \sum_{r=0}^{\infty} S_{r(\beta(k+1)+j)} \left(\frac{x-\mu}{\sigma}\right)^r \tag{13}$$

Where $S_{r(m)} = \sum_{k=R}^{\infty} (-1)^{k+R} \binom{m}{k} \binom{k}{r}$

Combining equations (12) and (13), the Weimal *cdf* can be expressed from equation (12) as:

$$F(X) = \sum_{r=0}^{\infty} \mathfrak{N}_r \phi\left(\frac{x-\mu}{\sigma}\right)^r = \sum_{r=0}^{\infty} \mathfrak{N}_r H(x; r, \mu, \sigma) \tag{14}$$

Where $\mathfrak{N}_r = \sum_{j,k=0}^{\infty} w_{j,k} S_{r(\beta(k+1)+j)}$

By differentiating equation (14) and changing indices, we can obtain the *pdf* of the Weimal distribution as:

$$f(x) = \frac{dF(x)}{dx} = \sum_{r=0}^{\infty} \mathfrak{N}_r r \phi\left(\frac{x-\mu}{\sigma}\right) \phi\left(\frac{x-\mu}{\sigma}\right)^{r-1} = \sum_{r=0}^{\infty} \mathfrak{N}_r h(x; r, \mu, \sigma) \tag{15}$$

Where $\mathfrak{N}_r = \sum_{j,k=0}^{\infty} w_{j,k} S_{r(\beta(k+1)+j)}$ and $\sum_{r=0}^{\infty} \mathfrak{N}_r = 1$

Equation (15) is the *pdf* of the Weimal distribution defined as a linear combination of EN *pdf*. So now, several properties of the Weimal distribution can be obtained by exploitation the properties of EN distribution.

3. Some Properties of the Weibull-Normal Distribution

3.1 Ordinary Moments

Moments are used to study some of the most important features and characteristics of a random variable such as mean (central tendency measure), variance (dispersion measure), skewness (Sk) and kurtosis (ku).

Let X_1, X_2, \dots, X_n denote a random sample from the standard Weibull-Normal (Weimal) distribution obtained from equation (14) for $\mu = 0$ and $\sigma = 0$, that is Weimal $(\alpha, \beta, 0, 1)$.

The n^{th} moment of X can be obtained as:

$$\mu_n = E(X^n) = \sum_{r=0}^{\infty} r \eta_r \int_{-\infty}^{\infty} X^n \phi(x) \phi(x)^{r-1} dx \tag{16}$$

Now, substituting for $\phi(x)$ and $\phi(x)^{r-1}$ in equation (16), using binomial expansion and simplifying, we have:

$$\mu_n = \sum_{r=0}^{\infty} \sum_{p=0}^{r-1} r \eta_r \frac{2^{-|r-1|}}{\sqrt{2\pi}} \binom{r-1}{p} / (n, p) \tag{17}$$

Where $/(n, p)$ represents the $(n, p)^{th}$ probability weighted moment (PWM) for any n and p positive integers of the standard normal distribution and is found as follows:

$$/(n, p) = \pi^{-\frac{p}{2}} 2^{\binom{n-1}{2}+p} \sum_{m_1=0}^{\infty} \dots \sum_{m_p=0}^{\infty} \frac{(-1)^{m_1+\dots+m_p}}{(2m_1+1)\dots(2m_p+1)m_1!\dots m_p!} X \Gamma\left(m_1 + \dots + m_p + \frac{p+n+1}{2}\right) \tag{18}$$

Now, according to Nadarajah [27],

$$(f)_n = \Gamma(f + n) / \Gamma(f) \tag{19}$$

And

$$F_{(A)}^{(n)}(a; b_1 \dots b_p; c_1 \dots c_p; x_1 \dots x_p) = \sum_{m_1=0}^{\infty} \dots \sum_{m_n=0}^{\infty} \frac{a_{m_1+\dots+m_p} (b_1)_{m_1} \dots (b_n)_{m_n} x_1^{m_1} \dots x_n^{m_n}}{(c_1)_{m_1} \dots (c_n)_{m_n} m_1! \dots m_n!} \tag{20}$$

Where equation (19) is the Lauricella function of type A [28],

Using these definitions in equation (19), equation (18) can be expressed as:

$$/(n, p) = \pi^{-\frac{p}{2}} 2^{\binom{n-1}{2}+p} F_{(A)}^{(p)}\left(\frac{p+n+1}{2}; \frac{1}{2}, \frac{3}{2}, \dots, \frac{3}{2}, -1, \dots, -1\right) \tag{21}$$

Given that $p + n$ is even.

Combining equations (17) and (21), it can be expressed that the n th moment of the standard Weimal distribution in terms of the Lauricella function of type A are demonstrated by Nadarajah, Lima *et al*, Pescim *et al*. [3, 6, 8, 9, 27]

$$\mu_n^l = \sum_{r=0}^{\infty} \sum_{p=0, p+n, even}^{r-1} \binom{r-1}{p} r \eta_r \pi^{-\frac{(p+1)}{2}} 2^{\frac{n}{2}-r+p} X \Gamma\left(\frac{p+n+1}{2}\right) F_{(A)}^{(p)}\left(\frac{p+n+1}{2}; \frac{1}{2}, \frac{3}{2}, \dots, \frac{3}{2}, -1, \dots, -1\right) \tag{22}$$

3.2 The Central Moments

The n th central moments or moment about the mean of X, say μ_n can be obtained as:

$$\mu_n = E[X - \mu_1^i]^n = \sum_{i=0}^n (-1)^i \mu_1^i \mu_{n-i}^i \tag{23}$$

The variance of X is the central moment of order two ($n=2$) and is given as:

$$Var(X) = \mu_2^i - (\mu_1^i)^2 \tag{24}$$

Where μ_2^i and $(\mu_1^i)^2$ are the second ordinary moment and first ordinary moment squared respectively which could be obtained using equation (22) as follows:

$$\mu_n^l = \sum_{r=0}^{\infty} \sum_{p=0, p+n, even}^{r-1} \binom{r-1}{p} r \eta_r \pi^{-\frac{(p+1)}{2}} 2^{\frac{n}{2}-r+p} X \Gamma\left(\frac{p+n+1}{2}\right) F_{(A)}^{(p)}\left(\frac{p+n+1}{2}; \frac{1}{2}, \frac{3}{2}, \dots, \frac{3}{2}, -1, \dots, -1\right) \tag{25}$$

For $n=1$, we therefore obtain the mean of the standard Weimal distribution from equation (22) as:

$$\mu_1^l = \sum_{r=0}^{\infty} \sum_{p=0, p+n, even}^{r-1} \binom{r-1}{p} r \eta_r \pi^{-\frac{(p+1)}{2}} 2^{\frac{1}{2}-r+p} X \Gamma\left(\frac{p+2}{2}\right) F_{(A)}^{(p)}\left(\frac{p+2}{2}; \frac{1}{2}, \frac{3}{2}, \dots, \frac{3}{2}, -1, \dots, -1\right) \tag{26}$$

Similarly, when $n=2$, we derive the following expression for the second moment as:

$$\mu_2^l = \sum_{r=0}^{\infty} \sum_{p=0, p+n, even}^{r-1} \binom{r-1}{p} r \eta_r \pi^{-\frac{(p+1)}{2}} 2^{1-r+p} X \Gamma\left(\frac{p+3}{2}\right) F_{(A)}^{(p)}\left(\frac{p+3}{2}; \frac{1}{2}, \frac{3}{2}, \dots, \frac{3}{2}, -1, \dots, -1\right) \tag{27}$$

Hence, the variance of $X \sim Weimal(\alpha, \beta, 0, 1)$ which is the second central moment of X is obtained as:

$$Var(X) = \mu_2^i - (\mu_1^i)^2 = \sigma^2 \tag{28}$$

Using equations (26) and (27) respectively.

The coefficient of skewness is the standardized third central moment of X about the mean and can be obtained using the expression:

$$Sk = E \left(\frac{X - \mu_1^I}{\sigma} \right)^3 = \frac{\mu_3}{\sigma^3} \tag{29}$$

Whereas the coefficient of kurtosis is the standardized fourth central moment of X about the mean and is given by

$$Ku = E \left(\frac{X - \mu_1^I}{\sigma} \right)^4 = \frac{\mu_4}{\sigma^4} \tag{30}$$

Where σ can be obtained using equation (28) while μ_3 and μ_4 are obtained using equation (23).

3.3 Moment generating function (mgf)

A general way of organizing all the moments into one mathematical object is called the mgf. In other words, the mgf generates the moments of X by differentiation, that is for any real number say k, the kth derivative of $M_x(t)$ evaluated at $t=0$ is the kth moment μ_k of X.

The mgf of a standard Weimal random variable X ($X \sim Weimal(\alpha, \beta, 0, 1)$) can be obtained as:

$$M_x(t) = E(e^{tx}) = \int_{-\infty}^{\infty} e^{tx} f(x) dx = \sum_{n=0}^{\infty} \frac{t^n}{n!} \mu_n^i \tag{31}$$

Where n and t are constants, t is a real number and μ_n^i denotes the nth ordinary moment of X and can be obtained from equation (22) as stated earlier.

$$M_x(t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \sum_{r=0}^{\infty} \sum_{p=0, p+n, even}^{r-1} \binom{r-1}{p} r \eta_r \pi^{-\frac{(p+1)}{2}} 2^{\frac{n}{2}-r+p} X \Gamma \left(\frac{p+n+1}{2} \right) F_{(A)}^{(p)} \left(\frac{p+n+1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{2}, -1, \dots, -1 \right) \tag{32}$$

3.4. Characteristics function (cf)

The characteristics function (cf) has many useful and important properties which give it a central role in statistical theory. Its approach is particularly useful in analysis of linear combination of random variables.

A representation for the cf is given by:

$$\phi_x(t) = E(e^{itx}) = E[\cos tx + i \sin tx] = E[\cos tx] + E[i \sin tx] \tag{33}$$

Simple algebra and power series expansion proves that:

$$\phi_x(t) = \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n}}{(2n)!} \mu_{2n}^i + \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!} \mu_{2n+1}^i \tag{34}$$

Where μ_{2n}^i and μ_{2n+1}^i are the moments of X for $n=2n$ and $n=2n+1$ respectively and can be obtained from μ_n^i in equation (25) as:

$$\mu_{2n}^i = \sum_{r=0}^{\infty} \sum_{p=0, p+n, even}^{r-1} \binom{r-1}{p} r \eta_r \pi^{-\frac{(p+1)}{2}} 2^{n-r+p} X \Gamma \left(\frac{p+2n+1}{2} \right) F_{(A)}^{(p)} \left(\frac{p+2n+1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{2}, -1, \dots, -1 \right) \tag{35}$$

And

$$\mu_{2n+1}^i = \sum_{r=0}^{\infty} \sum_{p=0, p+n, even}^{r-1} \binom{r-1}{p} r \eta_r \pi^{-\frac{(p+1)}{2}} 2^{n-r+p+0.5} X \Gamma \left(\frac{p+2n+2}{2} \right) F_{(A)}^{(p)} \left(\frac{p+2n+2}{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{2}, -1, \dots, -1 \right) \tag{36}$$

3.5 Order Statistics

Order statistics have been used in a wide range of problems including robust statistical estimation, detection of outliers, characterization of probability distribution, goodness of fit tests, entropy estimation, analyses of censored samples, reliability analysis, quality control and even researches bordering on strength of materials. In this section, closed form expression for the pdf's of the ith order statistics of the Weibull-Normal (that is, Weimal) distribution is derived.

Suppose X_1, X_2, \dots, X_n is a random sample from the standard Weimal distribution and let $X_{1:n}, X_{2:n}, \dots, X_{i:n}$ denote the corresponding order statistic obtained from this sample. The pdf, $f_{1:n}(x)$ of the ith order statistic can be obtained by:

$$F_x(x) = \frac{n!}{(i-1)!(n-i)!} f(x) F(x)^{i-1} [1 - F(x)]^{n-1} \tag{37}$$

Using equations (6) and (7), the pdf, $f_{1:n}(x)$ of the ith order statistic $X_{1:n}$ can be expressed from equation (37) as:

$$f_{1:n}(x) = \frac{n!}{(i-1)!(n-i)!} \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \left[\frac{\alpha \beta}{\sigma} \phi \left(\frac{x-\mu}{\sigma} \right) \frac{\phi \left(\frac{x-\mu}{\sigma} \right)^{\beta-1}}{\left[1 - \phi \left(\frac{x-\mu}{\sigma} \right) \right]^{\beta+1}} \exp \left\{ -\alpha \left[\frac{\phi \left(\frac{x-\mu}{\sigma} \right)}{1 - \phi \left(\frac{x-\mu}{\sigma} \right)} \right]^{\beta} \right\} \right] X \left[1 - \exp \left\{ -\alpha \left[\frac{\phi \left(\frac{x-\mu}{\sigma} \right)}{1 - \phi \left(\frac{x-\mu}{\sigma} \right)} \right]^{\beta} \right\} \right]^{i+k-1} \tag{38}$$

Hence the pdf of the minimum order statistic $X_{(1)}$ and maximum order statistic $X_{(n)}$ of the Weimal distribution respectively given by:

$$f_{1:n}(x) = n \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \left[\frac{\alpha\beta}{\sigma} \phi\left(\frac{x-\mu}{\sigma}\right) \frac{\phi\left(\frac{x-\mu}{\sigma}\right)^{\beta-1}}{\left[1-\phi\left(\frac{x-\mu}{\sigma}\right)\right]^{\beta+1}} \exp\left\{-\alpha \left[\frac{\phi\left(\frac{x-\mu}{\sigma}\right)}{1-\phi\left(\frac{x-\mu}{\sigma}\right)}\right]^\beta\right\} \right] X \left[1 - \exp\left\{-\alpha \left[\frac{\phi\left(\frac{x-\mu}{\sigma}\right)}{1-\phi\left(\frac{x-\mu}{\sigma}\right)}\right]^\beta\right\} \right]^k \tag{39}$$

And

$$f_{n:n}(x) = n \left[\frac{\alpha\beta}{\sigma} \phi\left(\frac{x-\mu}{\sigma}\right) \frac{\phi\left(\frac{x-\mu}{\sigma}\right)^{\beta-1}}{\left[1-\phi\left(\frac{x-\mu}{\sigma}\right)\right]^{\beta+1}} \exp\left\{-\alpha \left[\frac{\phi\left(\frac{x-\mu}{\sigma}\right)}{1-\phi\left(\frac{x-\mu}{\sigma}\right)}\right]^\beta\right\} \right] X \left[1 - \exp\left\{-\alpha \left[\frac{\phi\left(\frac{x-\mu}{\sigma}\right)}{1-\phi\left(\frac{x-\mu}{\sigma}\right)}\right]^\beta\right\} \right]^{n-1} \tag{40}$$

3.6. Estimation of Parameters of Weimal Distribution

The estimation of the parameters of the Weimal distribution is presented using the method of maximum likelihood in this section. Let X_1, X_2, \dots, X_n be a random sample from the Weimal distribution with unknown parameter $\theta = (\alpha, \beta, \mu, \sigma)^T$. The total log-likelihood function for θ is obtained from $f(x)$ as follows:

$$L(X_1, X_2, \dots, X_n / \alpha, \beta, \mu, \sigma) = \left(\frac{\alpha\beta}{\sigma}\right)^n \sum_{i=1}^n \left(\phi\left(\frac{x_i-\mu}{\sigma}\right)\right) \frac{\sum_{i=1}^n \phi\left(\frac{x_i-\mu}{\sigma}\right)^{\beta-1}}{\sum_{i=1}^n \left[1-\phi\left(\frac{x_i-\mu}{\sigma}\right)\right]^{\beta+1}} \exp \sum_{i=1}^n \left\{-\alpha \left[\frac{\phi\left(\frac{x_i-\mu}{\sigma}\right)}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right]^\beta\right\} \tag{41}$$

Let $l(\theta) = L(X_1, X_2, \dots, X_n / \alpha, \beta, \mu, \sigma)$ therefore

$$l(\theta) = n \log \alpha + n \log \beta - n \log \sigma - \frac{\pi}{2} \log(2\pi) - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 + (\beta - 1) \sum_{i=1}^n \log \left[\phi\left(\frac{x_i-\mu}{\sigma}\right)\right] - (\beta + 1) (\beta - 1) \sum_{i=1}^n \log \left[1 - \phi\left(\frac{x_i-\mu}{\sigma}\right)\right] - \alpha \sum_{i=1}^n \left[\frac{\phi\left(\frac{x_i-\mu}{\sigma}\right)}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right]^\beta \tag{42}$$

Meanwhile differentiating $l(\theta)$ partially with respect to each of the parameters: $\alpha, \beta,$ and σ and setting the results to zero gives the maximum likelihood estimates of the respective parameters. The partial derivative of $l(\theta)$ with respect to each parameter or the score function is given by:

$$U_n(\theta) = \left(\frac{\partial l\theta}{\partial \alpha}, \frac{\partial l\theta}{\partial \beta}, \frac{\partial l\theta}{\partial \mu}, \frac{\partial l\theta}{\partial \sigma}\right)$$

Where the components of the score vector $U(\theta) = (U_\alpha, U_\beta, U_\mu, U_\sigma)$ are:

$$U_\alpha = \frac{\partial l\theta}{\partial \alpha} = \frac{n}{\alpha} - \sum_{i=1}^n \left[\frac{\phi\left(\frac{x_i-\mu}{\sigma}\right)}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right]^\beta \tag{43}$$

$$U_\beta = \frac{\partial l\theta}{\partial \beta} = \frac{n}{\beta} + \sum_{i=1}^n \log \left[\frac{\phi\left(\frac{x_i-\mu}{\sigma}\right)}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right] - \alpha \sum_{i=1}^n \left[\frac{\phi\left(\frac{x_i-\mu}{\sigma}\right)}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right]^\beta \log \left[\frac{\phi\left(\frac{x_i-\mu}{\sigma}\right)}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right] \tag{44}$$

$$U_\mu = \frac{\partial l\theta}{\partial \mu} = \frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu) - \frac{(\beta-1)}{\sigma\sqrt{2\pi}} \sum_{i=1}^n \left\{\frac{\exp\left[-\frac{1}{2}\left(\frac{x_i-\mu}{\sigma}\right)^2\right]}{\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right\} - \frac{(\beta+1)}{\sigma\sqrt{2\pi}} \sum_{i=1}^n \left\{\frac{\exp\left[-\frac{1}{2}\left(\frac{x_i-\mu}{\sigma}\right)^2\right]}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right\} + \frac{\alpha\beta}{\sigma\sqrt{2\pi}} \sum_{i=1}^n \left[\frac{\phi\left(\frac{x_i-\mu}{\sigma}\right)}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right]^{\beta-1} \left\{\frac{\exp\left[-\frac{1}{2}\left(\frac{x_i-\mu}{\sigma}\right)^2\right]}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right\} \tag{45}$$

$$U_\sigma = \frac{\partial l\theta}{\partial \sigma} = -\frac{n}{\sigma} + \frac{1}{\sigma^3} \sum_{i=1}^n (x_i - \mu)^2 - \frac{(\beta-1)}{\sigma^2\sqrt{2\pi}} \sum_{i=1}^n \left\{\frac{(x_i-\mu)\exp\left[-\frac{1}{2}\left(\frac{x_i-\mu}{\sigma}\right)^2\right]}{\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right\} - \frac{(\beta+1)}{\sigma^2\sqrt{2\pi}} \sum_{i=1}^n \left\{\frac{(x_i-\mu)\exp\left[-\frac{1}{2}\left(\frac{x_i-\mu}{\sigma}\right)^2\right]}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right\} + \frac{\alpha\beta}{\sigma^2\sqrt{2\pi}} \sum_{i=1}^n \left[\frac{\phi\left(\frac{x_i-\mu}{\sigma}\right)}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right]^{\beta-1} \left\{\frac{(x_i-\mu)\exp\left[-\frac{1}{2}\left(\frac{x_i-\mu}{\sigma}\right)^2\right]}{1-\phi\left(\frac{x_i-\mu}{\sigma}\right)}\right\} \tag{46}$$

Maximization of equation (42) can be performed by using well established routines like `nim`-routine or `optimize` in the R-statistical package. Setting these equation to zero, (that is, $U(\theta) = 0$) and solving them simultaneously yields the maximum likelihood estimate (MLE) $\hat{\theta}$ of θ . These equations cannot be solved analytically and therefore statistical softwares can be used to solve them numerically by means of iterative techniques like Newton-Raphson method.

4. Conclusion

In this research article, a new-four-parameter probability model named Weibull-Normal distribution (also to be known as Weimal distribution) resulting from the hybridization of two well-known probability models, namely: Weibull distribution and Normal distribution, is introduced. The new probability model extends the classical normal distributions by adding skewness to it. An obvious reason for generalizing a classical distribution is the fact that the generalization provides more flexibility to analyze real-life data. The new distribution has proved to be versatile and analytically tractable during the generalization process. The Weimal identity function can be expressed as a linear combination of exponentiated normal density functions, thereby enabling

derivations of vital mathematical properties comprising of moments, moment generating function, characteristics function and order statistics. The estimation of the parameters has been approached by the method of maximum likelihood.

5. References

1. Azzalini A. A class of distributions which includes the normal ones. *Scandinavian Journal of Statistics*. 1985; 31:171-178.
2. Eugene N, Lee C, Famoye F. Beta-Normal distribution and its applications. *Communications in Statistics-Theory and Methods*. 2002; 31:497-512.
3. Nadarajah S. A generalized Normal distribution. *Journal of Applied Statistics*. 2005; 32:685-694.
4. Cordeiro GM, De Castro M. A new family of generalized distributions. *Journal of Statistical Computation and Simulation*. 2011; 81:883-989.
5. Cordeiro GM, Cintra RJ, R'Ego LC, Ortega EMM. The McDonald Normal distribution. *Pakistan journal of Statistics and Operation Research*. 2012; 8:301-329.
6. Nadarajah S, Kotz S. The exponentiated type distribution. *Acta Applicandae Mathematicae Amsterdam*. 2006; 92:97-111.
7. Alzaatreh A, Famoye F, Lee C. The Gamma Normal distribution: Properties and Applications. *Computational Statistics and Data Analysis*. 2014; 69:67-80.
8. Lima MS, Cordeiro GM, Ortega EMM. A new extension of the Normal distribution. *Journal of Data Science*. 2015; 13:385-408.
9. Pescim RR, Nadarajah S. The Kummer Beta Normal: A Skewed Model. *Journal of Data Science*. 2015; 13:509-532.
10. Shaw WT, Buckley LR. The alchemy of probability distributions beyond Gram-Charlier expansions and a skew-kurtotic normal distribution from a rank transmutation map, 2007. Available at: <http://library.wolfram.com/infocenter/Articles/6670/alchemy.pdf>
11. Zografos K, Balakrishnan N. On families of Beta- and generalized Gamma- generated distributions and associated inference. *Statistical Methods*. 2009; 63:344-362.
12. Alexander C, Cordeiro GM, Ortega EMM, Saraabia JM. Generalized Beta- generated distributions. *Computational Statistics and Data Analysis*. 2012; 56:1880-1897.
13. Ristic MM, Balakrishnan N. The Gamma-Exponentiated Exponential distribution. *Journal of Statistical Computation and Simulation*. 2012; 82:1191-1206.
14. Torabi H, Montazari NH. The Gamma-Uniform distribution and its application. *Kybernetika*. 2012; 48:16-30.
15. Amini M, Mirmostafaei SMTK, Ahmadi J. Log-Gamma generated families of distributions. *Statistics*. 2014; 48(4):913-932.
16. Alzaghal A, Lee C, Famoye F. Exponentiated T-X family of distributions with some applications. *International Journal of Probability and Statistics*. 2013; 2:31-49.
17. Cordeiro GM, Ortega EMM, da-Cunha DCC. The exponentiated generalized class of distribution. *Journal of Data Science*. 2013; 11:1-27.
18. Torabi H, Montazari NH. The Logistic-Uniform distribution and its application. *Communications in Statistics-Theory and Methods*. 2014; 43:2551-2569.
19. Alzaghal A, Famoye F, Lee C. Weibull-Pareto distribution and its applications. *Communications in Statistics-Theory and Methods*. 2013; 42(9):1673-1691.
20. Tahir MH, Cordeiro GM, Alzaatreh A, Mansoor M, Zubair M. The Logistic-X family of distributions and its applications. *Communications in Statistics-Theory and Methods*. 2016; 45(24):7326-7349.
21. Bourgeignon M, Silva RB, Cordeiro CGB. The Weibull-G family of probability distributions. *Journal of Data Science*. V; 12:53-68.
22. Alizadeh M, Cordeiro GM, De-Brito E, Demetrio CGB. The Beta Marshall-Olkin family of distributions. *Journal of Statistical Distribution and Applications*. 2015; 2(4):1-18.
23. Johnson NL, Kotz S, Balakrishnan N. *Continuous Univariate Distributions*, 2nd Edition, John Wiley, New York, 1994, 1.
24. Oguntunde PE, Balogun OS, Okagbue HI, Bishop SA. The Weibull-Exponential Distribution: Its Properties and Applications. *Journal of Applied Science*. 2015; 15(11):1305-1311.
25. Merovci F, Elbaral I. Weibull-Rayleigh Distribution: Theory and Applications. *Applied Mathematics and Information Sciences*. 2015; 9(5):1-11.
26. Afify MZ, Yousof HM, Cordeiro GM, Ortega EMM, Nofal ZM. The Weibull-Frechet distribution and its applications. *Journal of Applied Science*. 2016; 43(14):2608-2626.
27. Nadarajah S. Explicit expressions for the moments of order Statistics. *Statistics and Probability Letters*. 2008; 78(2):196-205.
28. Exton H. *Handbook of Hypergeometric integrals: Theory, Applications, Tables, Computer programs*. New York; Halsted Press, 1978.
29. Smith RL, Naylor JC. A comparison of maximum likelihood and Bayesian estimators for the three-parameter Weibull distribution. *Applied Statistics*. 1987; 36:358-369.
30. Barreto-Souza WM, Cordeiro GM, Simas AB. Some results for the Beta-Frechet distribution. *Communications in Statistics-Theory and Methods*. 2011; 40:798-811.
31. Afify AZ, Aryal G. The Kumaraswamy exponentiated Frechet distribution. *Journal of Data Science*. 2016; 6:1-19.
32. Nicholas MD, Padgen WJ. A bootstrap control chart for Weibull percentiles. *Quality Reliability Eng. Int*. 2006; 22:141-151.
33. Cordeiro GM, Lemonte AJ. The β -Birnbbaum-Saunders distribution: An improved distribution for fatigue life modeling. *Computational Statistics and Data Analysis*. 2011; 55:1445-1461.
34. Al-Aqtash R, Lee C, Famoye F. Gumbel-Weibull distribution: Properties and Applications. *Journal of Modern Applied Statistical Methods*. 2014; 13:201-225.
35. Afify AZ, Nofal WM, Butt NS. Transmuted complementary Weibull-Geometric distribution. *Pakistan Journal of Statistics and Operation Research*. 2014; 10:435-454.